

Marine Biodiversity Records

<http://journals.cambridge.org/MBD>

Additional services for **Marine Biodiversity Records**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



A new record of *Holothuria arguinensis* colonizing the Mediterranean Sea

Mercedes González-Wangüemert and Giomar Borrero-Pérez

Marine Biodiversity Records / Volume 5 / October 2012 / e105

DOI: 10.1017/S1755267212000887, Published online:

Link to this article: http://journals.cambridge.org/abstract_S1755267212000887

How to cite this article:

Mercedes González-Wangüemert and Giomar Borrero-Pérez (2012). A new record of *Holothuria arguinensis* colonizing the Mediterranean Sea. Marine Biodiversity Records, 5, e105 doi:10.1017/S1755267212000887

Request Permissions : [Click here](#)

A new record of *Holothuria arguinensis* colonizing the Mediterranean Sea

MERCEDES GONZÁLEZ-WANGÜEMERT¹ AND GIOMAR BORRERO-PÉREZ²

¹Centro de Ciências do Mar (CCMAR), CIMAR-Laboratório Associado, Universidade do Algarve, Gambelas, 8005-139 Faro, Portugal, ²Instituto de Investigaciones Marinas y Costeras 'José Benito Vives de Andrés' (INVEMAR), Cerro Punta Betín, AA 1016, Santa Marta, Colombia

Holothuria arguinensis is reported for the first time on the south-eastern Spanish coast. One specimen of this species was identified in El Mojón (Alicante, Spain, 37°50'54.17"N 0°45'39.90"W). This finding widens the geographical range of *H. arguinensis* and demonstrates that its environmental requirements are changing. Future genetic studies in this species would allow us to improve our understanding of its colonization from the Atlantic Ocean to the Mediterranean Sea.

Keywords: sea cucumbers, molecular barcoding, colonization, range expansion, water temperature, anthropogenic impact

Submitted 30 August 2012; accepted 7 September 2012

INTRODUCTION

The *Holothuria* genus is the only Holothuriidae present in the Mediterranean Sea and north-eastern Atlantic, being found in different marine habitats from coastal shallow waters. They are found in high densities providing important ecosystem services enhancing nutrient cycling and local productivity in oligotrophic carbonate sediments through their bioturbation and deposit feeding activities (Byrne *et al.*, 2010).

Holothuria (*Roweothuria*) *arguinensis* Koehler & Vaney, 1906, belongs to this genus and in recent years it is being considered as a target species in the increasing sea cucumber fishery (Aydin, 2008; Sicuro & Levine, 2011). This species had been considered to be a north-eastern Atlantic species that is distributed from Portugal to Morocco and Mauritania, including the Canary Islands (Thandar, 1988; Hanson, 2001). It has not been found in other Macaronesian Islands such as Açores, Selvagens or Madeira, or in the Cape Verde Archipelago (Pereira, 1997; Borrero-Pérez *et al.*, 2010; Micael *et al.*, 2012). However, its geographical distribution is changing, including the colonization of the Mediterranean Sea, where it was registered for the first time on the Granada coast, southern Spain (Ocaña & Pérez-Ruzafa, 2004) and its extension to the north Portugal coast in the Berlengas Islands (39°24.56'N 9°30.88'W) (Rodrigues, 2012).

Considering this framework, the aims of this paper are: (i) to record the occurrence for the first time of *H. arguinensis* to the north of the Alborán Sea (south-eastern Spain); (ii) to discuss the colonization process of this species; and (iii) to contribute to the clarification of the taxonomy of *Holothuria* genus species from the Atlanto-Mediterranean region.

MATERIALS AND METHODS

One specimen was collected in El Mojón (37°50'54.17"N 0°45'39.90"W), south-eastern Spanish coast. It was identified *in situ* through its external morphology and subsequently this identification was confirmed using traditional taxonomic characters, such as ossicles, on the basis of the original description of this species (Koehler & Vaney, 1906) and another relevant taxonomic paper (Thandar, 1988). In addition, cytochrome c oxidase I (COI) gene was amplified according to protocols from Borrero-Pérez *et al.* (2010) and this sequence was recorded in the GenBank database (accession number JX125700).

RESULTS

SYSTEMATICS

Class HOLOTHUROIDEA

Order ASPIDOCHIROTIDA

Family HOLOTHURIIDAE

Genus *Holothuria* Linnaeus, 1767

Sugenus *Roweothuria* Thandar, 1988

Holothuria (*Roweothuria*) *arguinensis* Koehler & Vaney, 1906 (Figures 1–3)

Holothuria arguinensis Koehler & Vaney, 1906, pp. 62–65, pl. 5, figures 5–13, pl. VI, figures 14–21. Cherbonnier, 1950, pp. 106–108, figure 3d.

Holothuria (*Lessonothuria*) *arguinensis* Rowe, 1969, pp. 149–150.

Holothuria (*Roweothuria*) *arguinensis* Thandar 1988, pp. 48 and 53.

Species length up to 185 mm; dorsal surface quite rounded and more or less wrinkled, cracked and evenly coloured brown on the entire region of bivium; on the sides, this colour is gone gradually to white; ventral surface slightly flattened, whitish and with punctuations that match the pedicels retracted which are light brown. Pedicels evenly distributed throughout

Corresponding author:

M. González-Wangüemert
Email: mwanguemert@ualg.pt




Fig. 1. Specimens of *Holothuria arguinensis* from localities belonging to its original geographical distribution. Individuals from Ria Formosa (Portugal) showing coloration variability: (A) dorsal side of individual AFM1; (B, C) dorsal and ventral sides of the individual AFM2. Ossicles of one individual (ACN1) from Canary Islands (Spain); (D) dorsal tables; (E) dorsal buttons; (F) ventral buttons. Scale bar: 20 μm.

the body, more numerous ventrally. Large papillae or warts on the dorsal surface, they are blackish brown to whitish and they are arranged in six longitudinal rows; two of these rows occupy the radial side and they constitute a whole by almost continuous rim; on bivium, each radius has two rows of papillae that alternated more or less irregularly with each other and

offer the same dimensions as the lateral papillae. Ossicles: tables few, mostly incomplete with four central openings and without marginal holes; rounded or angular edges; spire with four rods joined to one another by a transverse bar and ending in a cluster of teeth; 60–65 μm. Ventral buttons are oval, very thick, smooth, rounded edges and they vary in




Fig. 2. Specimen of *Holothuria arguinensis* collected in El Mojón (Spain): (A) complete specimen viewed from dorsal side; (B) complete specimen viewed from ventral side; (C) dorsal view of papilla; (D) ventral view of papilla; (E) dorsal buttons; (F) dorsal tables; (G) ventral buttons. Scale bar: 20 μm.




Fig. 3. Specimens of *Holothuria arguinensis*: (A, C, D) Ria Formosa (Portugal) during low tide (photographs: M. González-Wangüemert); (B) Arrábida Natural Park (Portugal) (photograph: E. Gonçalves).

size and number of holes, but the most common is 2–3 pairs of small holes; dorsal ones are flatter than the ventral face, and they are not quite the same form as those of the ventral which have rounded edges, while those of the dorsal surface are toothed, very rough, but it is possible to find all the transitions between these two extreme forms; 2–4 pairs of large holes; 50–100 μm . Numerous Polian vesicles; single stone canal (Koehler & Vaney, 1906; Thandar, 1988).

SPECIMEN FROM EL MOJÓN (SOUTH-EASTERN SPAIN)

On the basis of *H. arguinensis* specimens collected and studied previously by us from localities belonging to its original distribution, we must highlight the variability of the coloration on dorsal surface. It ranges from uniform brown to a coloration gradient (brown in the centre and white or yellowish on sides), but always with darker warts (Figure 1A–C). Another important observation is about the rough surface of ventral buttons: Koehler & Vaney (1906) were not clear about this characteristic in *H. arguinensis*; and Thandar (1988) described them as smooth buttons. However, they are rough although the dorsal buttons can be toothed too (Figure 1E, F).

The specimen collected at El Mojón (south-eastern Spain) fits with the characteristics of *H. arguinensis* presented previously. The length of specimen was 175 mm; dorsal surface is uniform brown with the darker warts alternating more or less irregularly (Figure 2A, B), and with the lateral large papillae arranged all along the continuous rim (Figure 2C, D). Dorsal and ventral ossicles fit with the description presented, observing on this specimen the rough surface of ventral buttons discussed previously (Figure 2E, G). According to the original description, the number of tables is low specially

in the ventral side showing incomplete discs, although some complete tables were also found with a marginal ring of holes and pointed edges (Figure 2F).

DISTRIBUTION AND HABITAT

Holothuria arguinensis is distributed in the north-east Atlantic, from the Berlengas Islands (Portugal) to Morocco and Mauritania, including the Canary Islands and the transition zone to the Mediterranean Sea (Alborán Sea). Its range of depth is from 0 to 52 m. *Holothuria arguinensis* is a species that does not take refuge in crevices, remains visible on sandy or seagrass beds, inhabiting the inter-tidal zone and being able to stand periods of drought during the low tides such as is observed in the Ria Formosa (Portugal) (Figure 3A–D).

TAXONOMIC REMARKS

The taxonomic status of the subgenus *Roweothuria*, *H. arguinensis* being the type species, should be revised. Three species *H. arguinensis*, *H. polii* and *H. vema* belong to this subgenus, considering mainly the spinose, flat rimmed table discs, and smooth to rough or spinose rosette-like buttons (Thandar, 1988). However, recent phylogenetic analysis showed that *H.(R.) arguinensis* is nearer to *H. (Holothuria) tubulosa*, *H. (Holothuria) mammata* and *H. (Vaneyothuria) lentiginosa* than *H.(R.) polii*, which is very divergent genetically (Borrero-Pérez *et al.*, 2010).

DNA BARCODING

The amplification of the COI mitochondrial gene, a sequence with 531 base pairs in length and the blast in GENBANK (<http://www.ncbi.nlm.nih.gov>), identified the individual from El Mojón as *H. arguinensis* with a 99% maximum identity with GQ214754.1 (Canary Islands, Spain) and GQ214755.1 (Algarve, Portugal) but showing a different haplotype which was recorded as JX125700.

DISCUSSION

The current new record of *H. arguinensis* is supported by the taxonomic information presented in this paper. In addition, this information will be useful to clarify the taxonomy of *Holothuria* genus species from the Atlanto-Mediterranean region, considering other records of this species in the Mediterranean Sea, which usually have been named as *Holothuria tubulosa* (Lorenzo-Heller, 2005). Therefore, this species could be under-sighted in the south of the Iberian Peninsula considering its similarity to other species.

The presence of *H. arguinensis* in El Mojón (south-eastern Spain) is probably related to climate changes and human impacts. The increase of temperatures in the north-eastern Atlantic Ocean (Brander *et al.*, 2003) would allow this species to adapt to warmer waters colonizing the Mediterranean Sea; this ability is reinforced by the high survival of this species to desiccation during the low tides (authors, personal observation in Ria Formosa and Olhos de Agua in South Portugal: Figure 3). Also, the increase of shipping traffic through the Strait of Gibraltar could be favouring the dispersal of this species, mainly during the settled pentaculata larvae (Asha & Muthiah, 2002; Ivy & Giraspy, 2006).

This finding widens the geographical range of *H. arguinensis* and demonstrates that its environmental requirements are

changing. Future genetic studies should allow us to improve our knowledge about the colonization events of this species from the Atlantic Ocean to the Mediterranean Sea.

ACKNOWLEDGEMENTS

This research was supported by CUMFISH project (PTDC/MAR/119363/2010; Principal Investigator: Dr M. González-Wangüemert) funded by Fundação para Ciência e Tecnologia (Portugal). M. González-Wangüemert was supported by a FCT postdoctoral fellowship (SFRH/BPD/70689/2010).

REFERENCES

- Asha P.S. and Muthiah P.** (2002) Spawning and larval rearing of sea cucumber *Holothuria (Theelothuria) spinifera* Theel. *SPC Beche-de-mer Information Bulletin* 16, 11–15.
- Aydin M.** (2008) The commercial sea cucumber fishery in Turkey. *SPC Beche-de-mer Information Bulletin* 28, 40–41.
- Borrero-Pérez G., Gómez-Zurita J., González-Wangüemert M., Marcos C. and Pérez-Ruzafa A.** (2010) Molecular systematics of the genus *Holothuria* in the Mediterranean and Northeastern Atlantic and a molecular clock for the diversification of the Holothuriidae (Echinodermata: Holothuroidea). *Molecular Phylogenetics and Evolution* 57, 899–906.
- Brander K., Blom G., Borges M.F., Erzini K., Henderson G., MacKenzie B.R., Mendes H., Ribeiro J., Santos A.M.P. and Torensen R.** (2003) Changes in fish distribution in the eastern North Atlantic: are we seeing a coherent response to changing temperature? *ICES Marine Science Symposia* 219, 261–270.
- Byrne M., Rowe F. and Uthicke S.** (2010) Molecular taxonomy, phylogeny and evolution in the family Stichopodidae (Aspidochirota: Holothuroidea) based on COI and 16S mitochondrial DNA. *Molecular Phylogenetics and Evolution* 56, 1068–1081.
- Hanson H.G.** (2001) Echinodermata. In Costello M.J., Emblow C. and White R. (eds) *European register of marine species: a checklist of the marine species in Europe and a bibliography of guides to their identification*. Collection Patrimoine Naturels 50. Paris: Muséum National d'Histoire Naturelle, 336–351.
- Ivy G. and Giraspy D.A.B.** (2006) Development of large-scale hatchery production techniques for the commercially important sea cucumber *Holothuria scabra* var. *versicolor* (Conand, 1986) in Queensland, Australia. *SPC Beche-de-mer Information Bulletin* 24, 28–38.
- Koehler R. and Vaney C.** (1906) Mission des Pêcheries de la Côte occidentale d'Afrique. II. Echinodermes. *Actes Société Linnéenne, Bordeaux* 60, 58–66, pls 4–6.
- Lorenzo-Heller M.A.** (2005) *Impacto bajo la cota cero: estudio sobre la interacción del buceo deportivo en las comunidades bentónicas del litoral granadino*. Master's thesis. University of Wales, Aberystwyth, 31 pp.
- Micael J., Alves M.J., Jones M.B. and Costa A.C.** (2012) Diversity of shallow-water asteroids (Echinodermata) in the Azorean Archipelago Research Articles. *Marine Biodiversity Records* 5. doi:10.17/S1755267211000534.
- Ocaña A. and Pérez-Ruzafa A.** (2004) Los equinodermos de las costas andaluzas. *Acta Granatense* 3, 83–136.
- Pereira M.O.R.** (1997) Checklist of the littoral echinoderms of the Azores. *Açoreana* 8, 331–337.
- Rodrigues N.** (2012) New geographic distribution records for Northeastern Atlantic species from Peniche and Berlengas Archipelago. *Arquipelago: Life and Marine Sciences* 29, 1–4.
- Rowe F.W.E.** (1969) A review of the family Holothuriidae (Holothuroidea: Aspidochirota). *Bulletin of the British Museum of Natural History (Zoology)* 18, 119–170.
- Sicuro B. and Levine J.** (2011) Sea cucumber in the Mediterranean: a potential species for aquaculture in the Mediterranean. *Reviews in Fisheries Science* 19, 299–304.
- and
- Thandar A.S.** (1988) A new subgenus of *Holothuria* with a description of a new species from the south-east Atlantic Ocean. *Journal of Zoology* 215, 47–54.

Correspondence should be addressed to:

M. González-Wangüemert
 Centro de Ciências do Mar (CCMAR)
 CIMAR-Laboratório Associado, Universidade do Algarve
 Gambelas, 8005-139 Faro, Portugal
 email: mwanguemert@ualg.pt